



Optical electron transfer through 2,7-diethynylfluorene spacers in mixed-valent complexes containing electron-rich " $(\eta^2\text{-dppe})(\eta^5\text{-C}_5\text{Me}_5)\text{Fe}$ " endgroups.

Floriane Malvoti, Cédric Rouxel, Olivier Mongin, Philippe Hapiot, Loïc Toupet, Mireille Blanchard-Desce, Frédéric Paul

► To cite this version:

Floriane Malvoti, Cédric Rouxel, Olivier Mongin, Philippe Hapiot, Loïc Toupet, et al.. Optical electron transfer through 2,7-diethynylfluorene spacers in mixed-valent complexes containing electron-rich " $(\eta^2\text{-dppe})(\eta^5\text{-C}_5\text{Me}_5)\text{Fe}$ " endgroups.. Dalton Transactions, 2011, 40 (25), pp.6616-8. 10.1039/c1dt10231a . hal-00842688

HAL Id: hal-00842688

<https://hal.science/hal-00842688>

Submitted on 15 Jul 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Optical electron transfer through 2,7-diethynylfluorene spacers in mixed-valent complexes containing electron-rich “ $(\eta^2\text{-dppe})(\eta^5\text{-C}_5\text{Me}_5)\text{Fe}^+$ ” endgroups†Floriane Malvolti,^a Cedric Rouxel,^b Olivier Mongin,^b Philippe Hapiot,^a Loic Toupet,^c Mireille Blanchard-Desce^b and Frédéric Paul^{*a}

Received 10th February 2011, Accepted 14th April 2011

DOI: 10.1039/c1dt10231a

We report in this communication the study of the intramolecular electron transfer through a 2,7-diethynylfluorenyl spacer in the Fe(II)/Fe(III) mixed-valent (MV) complex $[(\eta^2\text{-dppe})(\eta^5\text{-C}_5\text{Me}_5)\text{FeC}\equiv\text{C}(2,7\text{-C}_{21}\text{H}_{24})\text{C}\equiv\text{CFe}(\eta^5\text{-C}_5\text{Me}_5)(\eta^2\text{-dppe})][\text{PF}_6]$ (**1** [PF_6]). The complex is generated *in situ* by comproportionation from its homovalent dinuclear Fe(II) and Fe(III) parents (**1** and **1** [PF_6]₂). It is shown that electronic delocalization is much more effective through a 2,7-fluorenyl than through a 4,4'-biphenyl bridging unit.

In the field of molecular electronics,^{1–6} the understanding of structural features influencing the intramolecular electron transfer has become decisive for optimizing the design of smart molecular-sized devices.^{7,8} While electron-transfer through several (hetero)aromatic units has been thoroughly investigated,⁹ to our knowledge, no electronic coupling through 2,7-diethynylfluorenyl-based bridges has been reported so far. Indeed, among the various such dinuclear transition metal complexes reported,^{10,11} most of them featured electron-poor metal centers and presented no other stable and isolable redox state(s), except perhaps for a few derivatives reported by Raithby and Wong.^{12,13} But even in these cases, the properties of the mixed-valent state of these compounds were not investigated. In the frame of a project aimed at developing redox-switchable multiphotonic absorbers,^{14–17} we recently synthesized the homovalent Fe(II) and Fe(III) compounds **1** and **1** [PF_6]₂ (Chart 1) featuring the electron-rich “ $(\eta^2\text{-dppe})(\eta^5\text{-C}_5\text{Me}_5)\text{Fe}^+$ ” endgroups.

We thus logically wondered about the intramolecular electron-transfer properties of the Fe(II)/Fe(III) mixed valent (MV)





Chart 1

congener of these compounds (**1** [PF_6]).^{1,8} Likewise to its 4,4'-biphenyl analogue (**2** [PF_6]) previously studied in our group¹⁸ or to other related dinuclear carbon-rich MV derivatives,^{7,19–21} **1** [PF_6] certainly constitutes another example of organometallic molecular wire.^{8,9} In that respect, we have previously shown that a remarkably large electronic coupling ($H_{\text{FeFe}} \cong 145 \text{ cm}^{-1}$) was operative between the redox-active endgroups of **2** [PF_6] in spite of the large intramolecular distance between them (*ca.* 16 Å).¹⁸ The coplanarity of the fluorenyl unit, which should result in a better overlap of its π -manifold, is anticipated to increase the electronic coupling between the redox-active endgroups relative to a 4,4'-biphenyl spacer. However, the loss of axial symmetry with the 2,7-fluorene spacer might simultaneously result in detrimental quantum interferences.^{22,23} Thus, in order to clearly establish how the electronic coupling is modified when progressing from **1** [PF_6] to **2** [PF_6], we have decided to experimentally determine H_{FeFe} for **1** [PF_6].

Given that a MV complex is always in equilibrium with its homovalent parents, we first wondered about the thermodynamic stability of **1** [PF_6] in the comproportionation equilibrium (Scheme 1).²⁴ Based on the well known eqn (1),²⁵ it was immediately apparent from the cyclic voltammogram of **1** (or **1** [PF_6]₂) that



Scheme 1

^aSciences Chimiques de Rennes, UMR CNRS 6226, Université de Rennes 1, Campus de Beaulieu, 35042, Rennes Cedex, France. E-mail: frederic.paul@univ-rennes1.fr

^bChimie et Photonique Moléculaires, UMR CNRS 6510, Université de Rennes 1, Campus de Beaulieu, 35042, Rennes Cedex, France

^cInstitut de Physique de Rennes, UMR CNRS 6251, Université de Rennes 1, Campus de Beaulieu, 35042, Rennes Cedex, France

† Electronic supplementary information (ESI) available: Synthesis of **1** and **1** [PF_6]₂. CIF file for **1**. Simulation procedure of the voltammogram used for **1**. UV and IR spectra obtained from equimolar mixtures of **1** and **1** [PF_6]₂. Solvatochromy and deconvolution of the near-IR absorption. CCDC reference number 694575. For crystallographic data in CIF or other electronic format see DOI: 10.1039/c1dt10231a

the corresponding MV compound has not a sufficient stability to be quantitatively isolated by chemical oxidation of **1**, since the voltammogram exhibits strongly overlapping waves for the two stepwise (and chemically reversible) iron-centred oxidations (see ESI†). Moreover, when coming close together, the apparent potential difference between these redox waves does not match anymore with the real ΔE° value.²⁶ Accurate values of the corresponding redox potentials in dichloromethane were obtained by simulation of this voltammogram (−0.11 and −0.22 V vs. ECS) and a thermodynamic equilibrium constant (K_c) of 76 ± 8 could be derived for this comproportionation reaction at 25 °C from the difference between these redox potentials ($\Delta E^\circ = 111 \pm 3$ mV).

$$(RT/F)\log(K_c) = \Delta E^\circ \quad (1)$$

Notably, this potential separation is significantly larger than that previously determined for **2**[PF₆] (62 mV), in line with a larger electronic interaction taking place between the redox endgroups in **1**[PF₆], which also translates in a larger stability constant for this MV complex (a K_c value of 11 had been found for **2**[PF₆]). While ensuring a good solubility of the compound, the two butyl chains on the fluorene spacer possibly contribute to increase ΔE° by limiting any pi-stacking or ion pairing in solution (Fig. 1). Interestingly, when the cyclic voltammetry is repeated in acetone, the separation between the redox waves is hardly discernible, indicating a much smaller comproportionation constant in this solvent possessing a higher dielectric constant. Under these conditions, simulation is needed to derive ΔE° and a K_c value of 28 ± 6 can be found for **1**[PF₆] in acetone.




Fig. 1 ORTEP of **1** (50% probability level).

In order to study **1**[PF₆] in CH₂Cl₂, the mixed valent complex was generated after solubilizing equimolar quantities of **1** and **1**[PF₆]₂ in this solvent.^{27,28} According to the K_c value found, **1**[PF₆] will be present in 81% in solution along with 9.5% of **1** and 9.5% of **1**[PF₆]₂. The infrared spectrum of this medium in the 1900–2000 cm^{−1} range reveals the presence of two new strong $\nu_{\text{Fe}\equiv\text{C}}$'s overlapping the corresponding modes of **1** and **1**[PF₆]₂ (ESI†). These absorptions, typical of Fe(II)–C≡C and a Fe(III)–C≡C acetylides, evidence the presence of distinct and localized redox sites in **1**[PF₆].²⁹ Also, after thawing an equimolar mixture of **1** and **1**[PF₆]₂ in CH₂Cl₂/1,2-C₂H₂Cl₂, a rhombic signal can be detected by ESR at 77 K ($g_1 = 1.983$, $g_2 = 2.035$ and $g_3 = 2.380$). The latter is typical of a low-spin Fe(III)-centred radical in cationic piano-stool acetylide complexes.³⁰ In accordance with the voltammetric study, these data establish **1**[PF₆] as a Class-II MV compound in the classification of Robin and Day.^{31,32}

We next turned our attention to the electronic absorptions of **1**[PF₆]. In the UV-vis range, the equimolar mixture between **1**

and **1**[PF₆]₂ reveals a spectrum resembling that of the parent homovalent complexes. However, several among the most intense transitions are shifted relative to these detected for pure **1** and **1**[PF₆]₂, in line with the presence of **1**[PF₆] in the medium. Then, in the near-IR range, an intense and *new* absorption is detected at ca. 5075 cm^{−1} which was absent for pure solutions of **1** and **1**[PF₆]₂. Actually, **1** is silent in this spectral range, whereas **1**[PF₆]₂ presents only a very weak absorption at 5423 cm^{−1} ($\epsilon = 410 \text{ M}^{-1} \text{ cm}^{-1}$) corresponding to a forbidden ligand field (LF) transition.^{18,30} The broad absorption at 2000 nm is thus diagnostic of the MV complex **1**[PF₆]. Considering the energy ($\tilde{\nu}_{\text{max}} = 4995 \text{ cm}^{-1}$) of the maximum, the halfwidth [$(\tilde{\nu}_{1/2})_{\text{exp}} > 6000 \text{ cm}^{-1}$] and also the (non-gaussian) shape of this band (Fig. 2), it cannot correspond to a single intervalence charge transfer (IVCT) band (see eqn (2)) but most likely results from several overlapping sub-bands.




Fig. 2 Near-IR spectra for an equimolar mixture of **1** and **1**[PF₆]₂ in dichloromethane ($1.3 \times 10^{-4} \text{ M}$ in total) and proposed de-convolution (sub-bands A–E: see text). Apparent extinction coefficients ($0.81 \times \epsilon$) are given in ordinate.

The near-IR band was then deconvoluted into several Gaussian sub-bands (ESI†). One evident contribution (A) corresponds to the weak ligand LF transition of **1**[PF₆]₂ present in the medium (9.5%), but its impact on the band shape is negligible. Another contribution (B) certainly originates from a similar LF transition taking place on the Fe(III) site of **1**[PF₆].³³ Again this sub-band should not be strongly influencing the band shape of the near-IR transition detected for the MV complex. Actually, (at least) three much more intense contributions need to be considered to properly reproduce the observed band shape (C–E). Notably, three such sub-bands were often considered in near-IR band deconvolutions for related organoiron MV derivatives.^{18,34,35} For MV complexes possessing an unpaired electron on metal d sub-levels lying close in energy to the HOMO-1/HOMO-2 levels, in pseudo-octahedral environments, when a sizable metal–metal interaction is mediated by the bridging ligand, these might be associated with the sought IVCT and two interconfigurational transitions.³⁶ Thus, similarly to what we did with the MV complex (η^2 -dppe)(η^5 -C₅Me₅)FeC≡C(1,4-C₆H₄)C≡CF(η^5 -C₅Me₅)(η^2 -dppe) (**3**[PF₆]),^{18,30} the most intense sub-band at lowest energy (C) was considered to correspond to the IVCT transition. In line with this hypothesis, this sub-band (along with D and E) is hypsochromically shifted when the dichloromethane solvent is changed for a solvent with a higher dielectric constant such as acetone or acetonitrile. This observation allows us to refine further the classification of the **1**[PF₆] as a Class-IIA MV complex.

$$(\tilde{\nu}_{1/2})_{\text{theo}} = (2310 \cdot \tilde{\nu}_{\text{max}})^{1/2} \quad (2)$$

$$H_{\text{ab}} = (2.06 \times 10^{-2} / d_{\text{ab}}) (\epsilon_{\text{max}} \tilde{\nu}_{\text{max}} \Delta \tilde{\nu}_{1/2})^{1/2} \quad (3)$$

The sub-bands C–E were considered to have similar halfwidths in the deconvolution procedure,³¹ with $\tilde{\nu}_{1/2}$ constrained to remain close to the theoretical value predicted by the Hush theory (eqn (2)).^{37,38} A fair fit can be obtained by this approach (Fig. 2), yielding a sensible reorganisation energy for the IVCT ($\lambda \approx 4670 \text{ cm}^{-1}$; $\tilde{\nu}_{1/2} = 3050 \text{ cm}^{-1}$; $\epsilon = 6050 \text{ M}^{-1} \text{ cm}^{-1}$) and sensible transition energies for the two interconfigurational processes D and E (7140 cm^{-1} and 10100 cm^{-1}).¹⁸

Finally, using the classic Hush equation (eqn (3)), a H_{FeFe} value of *ca.* 380 cm^{-1} was derived for **1**[PF₆] from the energy, intensity and halfwidth of the sub-band C (IVCT). These electronic couplings are quite important for two metal centres located *ca.* 16 \AA apart (Fig. 1).^{9,39} This nanoscopic distance outlines further the strong delocalization of the unpaired electron–hole in this organometallic MV compound. In line with established trends,^{25,40} the larger H_{FeFe} value found for **1**[PF₆] relative to **2**[PF₆], confirms our initial belief that the co-planarity of the central aromatic spacer is beneficial to the electronic communication. This rather large H_{FeFe} value found for **1**[PF₆], along with the intensity of its IVCT absorption, is reminiscent of signatures usually observed for more strongly coupled MV complexes (Class-IIB), such as **3**[PF₆] previously mentioned.¹⁸

In conclusion, we have established here that the 2,7-fluorenyl spacer allows us to significantly improve the electronic communication between the $[(\eta^2\text{-dppe})(\eta^5\text{-C}_5\text{Me}_5)\text{Fe}]^{n+}$ ($n = 0, 1$) redox-active endgroups compared with the 4,4'-diethynylbiphenyl spacer in the corresponding MV complexes. In dichloromethane, when compared with **2**[PF₆], this translates in a large increase of the comproportionation constant (K_c) of **1**[PF₆] (Scheme 1) and in a smaller improvement (2.6 fold) of the electronic coupling between the redox endgroups. However, according to the H_{FeFe} value(s) found, the end-to-end electron transfer is now clearly adiabatic for **1**[PF₆] ($H_{\text{FeFe}} \geq 200 \text{ cm}^{-1}$),⁴¹ and becomes much faster than in the Class-IIA MV analogue **2**[PF₆].

Acknowledgements

CNRS and UEB (EPT project) are acknowledged for specific financial supports.

Notes and references

- 1 M. Akita and T. Koike, *J. Chem. Soc., Dalton Trans.*, 2008, 3523–3530.
- 2 P. J. Low, *J. Chem. Soc., Dalton Trans.*, 2005, 2821–2824.
- 3 N. Robertson and G. A. Mc. Gowan, *Chem. Soc. Rev.*, 2003, **32**, 96–103.
- 4 R. L. Carroll and C. B. Gorman, *Angew. Chem., Int. Ed.*, 2002, **41**, 4379–4400.
- 5 J.-P. Launay and C. Coudret, in *Electron Transfer in Chemistry*, ed. V. Balzani and A. P. de Silva, Wiley-VCH, 2000, vol. 5, pp. 3–47.
- 6 J. M. Tour, *Acc. Chem. Res.*, 2000, **33**, 791–803.
- 7 T. Ren, *Organometallics*, 2005, **24**, 4854–4870.

- 8 F. Paul and C. Lapinte, *Coord. Chem. Rev.*, 1998, **178–180**, 431–509.
- 9 (a) J.-P. Launay, *Chem. Soc. Rev.*, 2001, **30**, 386–397; (b) S. F. Nelsen, *Chem.–Eur. J.*, 2000, **4**, 581–588.
- 10 W.-Y. Wong and C.-L. Ho, *Coord. Chem. Rev.*, 2006, **250**, 2627–2690.
- 11 W.-Y. Wong, *Coord. Chem. Rev.*, 2005, **249**, 971–997.
- 12 J. Lewis, P. R. Raithby and W.-Y. Wong, *J. Organomet. Chem.*, 1998, **556**, 219–228.
- 13 W.-Y. Wong, G.-L. Lu, K.-F. Ng, C.-K. Wong and K.-H. Choi, *J. Organomet. Chem.*, 2001, **637–639**, 159–166.
- 14 M. P. Cifuentes, M. G. Humphrey, J. P. Morrall, M. Samoc, F. Paul, T. Roisnel and C. Lapinte, *Organometallics*, 2005, **24**, 4280–4288.
- 15 O. Mongin, L. Porrès, M. Charlot, C. Katan and M. Blanchard-Desce, *Chem.–Eur. J.*, 2007, **13**, 1481–1498.
- 16 N. Gauthier, C. Olivier, S. Rigaut, D. Touchard, T. Roisnel, M. G. Humphrey and F. Paul, *Organometallics*, 2008, **27**, 1063–1072.
- 17 N. Gauthier, G. Argouarch, F. Paul, L. Toupet, A. Ladjarafi, K. Costuas, J.-F. Halet, M. Samoc, M. P. Cifuentes, T. C. Corkery and M. G. Humphrey, *Chem.–Eur. J.*, 2011, **17**, 5561–5577.
- 18 S. Ibn Ghazala, F. Paul, L. Toupet, T. Roisnel, P. Hapiot and C. Lapinte, *J. Am. Chem. Soc.*, 2006, **128**, 2463–2476.
- 19 S. Rigaut, D. Touchard and P. H. Dixneuf, in *Redox Systems Under Nano-space Control*, ed. T. Hirao, Springer Verlag, Heidelberg, 2006.
- 20 F. de Montigny, G. Argouarch, K. Costuas, J.-F. Halet, T. Roisnel, L. Toupet and C. Lapinte, *Organometallics*, 2005, **24**, 4558–4572.
- 21 T. Weyland, K. Costuas, L. Toupet, J.-F. Halet and C. Lapinte, *Organometallics*, 2000, **19**, 4228–4239.
- 22 V. Marvaud, J.-P. Launay and C. Joachim, *Chem. Phys.*, 1993, **177**, 23–30.
- 23 C. Patoux, C. Coudret, J.-P. Launay, C. Joachim and A. Gourdon, *Inorg. Chem.*, 1997, **36**, 5037–5049.
- 24 S. Szafert, F. Paul, W. E. Meyer, J. A. Gladysz and C. Lapinte, *C. R. Chim.*, 2007, **10**, 693–701.
- 25 D. Astruc, *Electron Transfer and Radical Processes in Transition-Metal Chemistry*, VCH Publishers, Inc., New-York Weinheim Cambridge, 1995.
- 26 D. E. Richardson and H. Taube, *Inorg. Chem.*, 1981, **20**, 1278–1285.
- 27 F. Paul, *work in progress*.
- 28 The syntheses of the homovalent complexes are given in the Supporting Information†.
- 29 F. Paul, J.-Y. Mevellec and C. Lapinte, *J. Chem. Soc., Dalton Trans.*, 2002, 1783–1790.
- 30 F. Paul, L. Toupet, J.-Y. Thépot, K. Costuas, J.-F. Halet and C. Lapinte, *Organometallics*, 2005, **24**, 5464–5478.
- 31 B. S. Brunshwig, C. Creutz and N. Sutin, *Chem. Soc. Rev.*, 2002, **31**, 168–184.
- 32 M. B. Robin and P. Day, *Adv. Inorg. Chem. Radiochem.*, 1967, **247–422**.
- 33 During the deconvolution process, this weak contribution was thus simulated with a comparable halfwidth than sub-band A and its energy was kept close to that of A.
- 34 M.-I. Chung, X. Gu, B. A. Etzenhouser, A. M. Spuches, P. T. Rye, S. K. Seetharaman, D. J. Rose, J. Zubietta and M. B. Spangler, *Organometallics*, 2003, **22**, 3485–3494.
- 35 Y. Matsuura, Y. Tanaka and M. Akita, *J. Organomet. Chem.*, 2009, **694**, 1840–1847.
- 36 *i.e.* IVCT transitions involving lower lying d electrons than these of the very frontier MOs, see for instance: K. D. Demadis, C. M. Hartshorn and T. J. Meyer, *Chem. Rev.*, 2001, **101**, 2655–2685.
- 37 G. C. Allen and N. S. Hush, *Prog. Inorg. Chem.*, 1967, **8**, 357–389.
- 38 N. S. Hush, *Prog. Inorg. Chem.*, 1967, **8**, 391–444.
- 39 **Crystal Data**. C₉₇H₁₀₂Fe₂P₄C₆H₄; $M = 1581.47$; Orthorhombic; $a = 20.690(2) \text{ \AA}$, $b = 20.374(2) \text{ \AA}$, $c = 40.538(3) \text{ \AA}$; $U = 17088(3) \text{ \AA}^3$, $T = 110 \text{ K}$; space group Pbc_a; $Z = 8$, 182259 reflection measured, 18444 unique ($R_{\text{int}} = 0.1363$) which were used in calculations. The final $wR(F_2)$ was 0.1275 (all data). CCDC deposition N^o: CCDC 694575.
- 40 S. Woitellier, J.-P. Launay and C. Joachim, *Chem. Phys.*, 1989, **131**, 481–488.
- 41 M. N. Paddon-Row, *Adv. Phys. Org. Chem.*, 2003, **38**, 1–85.